

【No.81】ノギテカン(トポテカン)療法

病名 _____ 外来開始日 _____ 年 月 日 治療開始日 _____ 年 月 日
 変更日 ① _____ 年 月 日 変更日 ② _____ 年 月 日
 _____ 科 ID _____ 氏名 _____ (才)

身長 _____ cm 体重 _____ kg 体表面積 _____ m²

主治医 _____ 指示医 _____ 確認医(科長) _____ PS 0・1・2・3・4

HBs抗原(+)→HBV DNA()、HBs抗原(-)→HBc抗体() HBs抗体()

指示内容		D1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21		
		月日																						
		曜日																						
① ○ポート患者 生食20mlシリンジ(穿刺) 生食250mlでルート確保後、ロック ○末梢患者 生食250mlでルート確保後、ロック	指示	●	●	●	●	●																		
	受領																							
	入力																							
	実施																							
	みどり																							
② {グラニセトンバッグ100ml or 生食100ml} +デキサート _____mg+ポララミン __A +ファモチジン __A (30min)	指示	●	●	●	●	●																		
	受領																							
	入力																							
	実施																							
	あか																							
③ 生食100ml +ハイカムチン _____mg (1.25~1.5mg/m ²) (1.1mg __V) (30min)	指示	●	●	●	●	●																		
	受領																							
	入力																							
	実施																							
	あか																							
④ ①の残液でフラッシュ	指示	●	●	●	●	●																		
	受領																							
	入力																							
	実施																							
⑤ 生食20ml シリンジロック or ヘパリンシリンジロック (ポート患者のみ)	指示	●	●	●	●	●																		
	受領																							
	入力																							
	実施																							

嘔吐リスク:軽度
(10~30%)

3週毎