

No.126 CPT-11/CDDP療法

病名 _____ 外来開始日 _____ 年 月 日 治療開始日 _____ 年 月 日
 変更日 ① _____ 年 月 日 変更日 ② _____ 年 月 日
 _____ 科 ID _____ 氏名 _____ (才)
 身長 _____ cm 体重 _____ kg 体表面積 _____ m²
 主治医 _____ 指示医 _____ 確認医(科長) _____ PS 0・1・2・3・4

HBs抗原(+) \rightarrow HBV DNA() ,HBs抗原(-) \rightarrow HBc抗体() HBs抗体()

指示内容	月日	D1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
		曜日																											
① ○ポート患者 生食20mlシリンジ(穿刺) 生食250mlでルート確保後、ロック ○末梢患者 生食250mlでルート確保後、ロック	指示								●																				
	受領	●																											
	入力																												
	実施																												
② 生食100ml+アロキシ(0.75) 1 A +デキサート 9.9mg +プロイメンド150mg1A (30min)	指示																												
	受領	●																											
	入力																												
	実施																												
②' 生食100ml+アロキシ(0.75) 1 A +デキサート 9.9mg (30min)	指示																												
	受領																												
	入力																												
	実施																												
③ 生食250ml+イリノテカン _____mg (60mg/m ²) (100mg _____V,40mg _____V) (60min) ※day8,15は終了後①でフラッシュ ※UGT1A1の測定について、事前に検討すること。	指示																												
	受領	●																											
	入力																												
	実施																												
④ KN1号 500ml + 硫酸Mg補正液20ml 0.4A +アスパラK注10ml 1A (60min)	指示																												
	受領	●																											
	入力																												
	実施																												
⑤ 20%マンニトール注 300ml 0.66V (30min)	指示																												
	受領																												
	入力																												
	実施																												
⑥ 生食500ml 1B +シスプラチン _____mg (60mg/m ²) (50mg _____V,25mg _____V,10mg _____V)(60min)	指示																												
	受領	●																											
	入力																												
	実施																												
⑦ YDソリタT3 500ml (60min)	指示																												
	受領	●																											
	入力																												
	実施																												
⑧ 生食20ml シリンジロック or ヘパリンシリンジロック (ポート患者のみ)	指示																												
	受領	●																											
	入力																												
	実施																												
内服 デカドロン8mg/body day2-4	指示																												
	受領		●	●	●																								
	入力																												
	実施																												

嘔吐リスク:中等度 (30~90%)

嘔吐リスク:高度 (>90%)